

CR

CASE REPORT

E. P. C. M. S.

Restoration of Elbow Flexion for a C5-C6 Brachial Plexus Injury Using a Double Nerve Transfer

Adham Abdelfattah, B.S.
Miguel Pirela-Cruz, M.D.

BACKGROUND INFORMATION

Many peripheral nerve injuries were previously repaired by direct means at the site of injury.¹ In contrast, the concept of nerve transfers to address peripheral nerve injuries that are not repairable by direct means is relatively new. In general, nerve injuries that have a long distance to regenerate to their respect target (that usually fail) or have a poor prognosis for recovery are lesions that may be considered for nerve transfers. This case demonstrates the concept of peripheral nerve transfers to reinnervate the biceps and brachialis using less critical nerves to restore elbow flexion.

CASE REPORT

A 28-year-old right-hand dominant male was involved in a motorcycle accident and sustained multiple injuries which included: C7-T1 cervicothoracic subluxation with contusion of the spinal cord and associated left lower extremity weakness, multiple rib fractures, pulmonary contusions, a grade III AC separation of the left shoulder and a left C5-C6 brachial plexus injury. In his evaluation, an MRI was performed with findings of bilateral C7 and T1 facet dislocation. He subsequently underwent cervicothoracic fusion to stabilize the spine from C6 to T2 vertebra for vertebral injuries associated with a left lower extremity neurologic impairment. In addition to these findings, he was found to have concomitant left C5-C6 brachial plexopathy with the typical findings of absence of active elbow flexors and dynamic shoulder stabilizers.

Approximately one year post-motorcycle accident, nerve conduction and needle electromyographic studies clearly demonstrated the affected muscles including those following the superior trunk of left brachial plexus: infraspinatus, supraspinatus, biceps, brachioradialis, teres major, teres minor, and deltoid muscles. This study also found absence of sensory conduction in the thumb indicating lesion distal to dorsal root ganglion and absence of denervation of rhomboid muscles indicating lesion distal to branching of dorsal scapular nerve. About 1 year 4 ½ months later, operative interventions for restoration of upper extremity functions via nerve transfers was undertaken.

Surgical approach of the peripheral nerves was carried out using a medial approach. The median and ulnar nerves were identified in the arm (figure 1.). Vessel loops were placed about the nerves for gentle retraction and further dissection was made for proximal

exposure of the nerves. Careful and meticulous dissection to free the ulnar and median nerves was used. A disposable nerve stimulator was used intraoperatively to identify the portion(s) of the nerves suitable for transfer. The nerve fibers located on the posteromedial aspect of the ulnar nerve which innervates the flexor carpi ulnaris was chosen to transfer. On the median nerve, fibers located on the anterolateral portion of the nerve which controls the FDS to the digits was determined suitable transfer.



Figure 1: Exposure of left brachial plexus

Next, identification of biceps branch of the musculocutaneous nerve was undertaken and secured with 9-0 Nylon suture, fibrin glue and a Neurogen wrap in an end-to-end repair to the ulnar branch of the FCU. Then an end-to-end repair of the median nerve fibers to the brachialis was performed. The wound was closed in a layered fashion using 2-0 Vicryl for the deep tissues and staples for the skin. A sterile soft dressing, followed by a long-arm posterior splint was placed.

OUTCOME

Patient has been followed for twelve months status post double nerve transfer of the left upper extremity brachial plexus. There were no pre-, intra-, or post-operative complications. Incision healed successfully as seen in Figure 2.

Continued on page 16



Figure 2: Left upper extremity

At approximately nine months post-operatively he was able to flex his left elbow (Figure 3). The patient was very satisfied with his elbow function, which he knew would get stronger over time.

At approximately nine months post-operatively he was able to flex his left elbow (Figure 3). The patient was very satisfied with his elbow function, which he knew would get stronger over time.



Figure 3: Left Elbow Flexion

Further post-operative testing demonstrated the ulnar and median nerves function to be essentially intact despite the fibers taken from these nerves for the transfers. The patient demonstrated very good hand function (Figure 4, 5 & 6).



Figure 4: Left-Hand: Finger Abduction



Figure 5: Left Hand: Fist



Figure 6: Wrist extension and finger flexion

DISCUSSION

Devastating and debilitating brachial plexus injuries can render dramatic changes in the lives of these patients to loss of upper extremity functions. Management is holistic, multidisciplinary and extensive in order to return the patient back to- or close to their baseline functions in society. Initial evaluation includes: a thorough history, physical examination with adjunctive imaging and electrophysiological studies. Primary goal in treatment should create efforts to return and maintain as much function of the upper extremity as possible. It is important to consider timing of restoration procedure implemented due to irreversible changes to the motor endplate and intraneural fibrosis that produces muscle resistance to neuronal stimuli that may occur 18 to 24 months post-injury negating use of nerve reconstruction and late type interventions.⁹ Early interventions include procedures in nerve repair or reconstruction using direct repair, neurolysis for scar compression, nerve grafting, and nerve transfers; or late interventions consisting of tendon transfers or pedicled muscle transfers of functioning muscles (ie, bipolar pectoralis muscle transfer or latissimus dorsi muscle transfer) to maintain shoulder function and stability along with elbow flexion.⁹

Successful nerve transfers to improve elbow flexion first began with Oberlin as a single nerve transfer to innervate the biceps.³⁻⁵ Unfortunately, some patients did not achieve the elbow flexion strength required for activities of daily living with this transfer. Later, the Oberlin transfer was modified by Mackinnon by adding

Continued on page 17

a second nerve transfer to reinnervate the brachialis muscle achieving double nerve transfers.⁶⁻⁸ Thereafter, many studies undertaken have provided supportive evidence in promising nerve transfers in returning shoulder function, elbow flexion, and sensation to upper extremities using a variety of methods and donor/recipient nerves for transfers.^{3-8,10,13-16} As in this patient, this mode of repair has improved elbow flexion. Moreover, double nerve transfers have been indicated above single transfers in achieving better strength in flexion.¹² Nerve transfers have offered benefits above nerve coaptation in proximal nerve lesions² which risks tension due to approximation in the cords increasing gap defect and/or elongated timing to repair that promotes risk of nerve retraction, also contributing to gap defect.¹ Additional nerve transfers that have been described to augment Oberlin-Mackinnon transfer include: the distal accessory nerve to the suprascapular nerve transfer and the branch to the lateral head of the triceps to the anterior branch of the axillary nerve to help provide reinnervation to the shoulder stabilizers.⁹

Nerve transfers offer a new approach to difficult peripheral nerve injuries. The double nerve transfer to reinnervate the biceps and brachialis is a reasonable approach for a C5-C6 brachial plexus injury and is now considered the treatment of choice for this lesion. Other nerve transfers are currently being developed to restore muscle and sensory function using less critical or expendable nerves.

REFERENCES

1. Trumble TE, McCallister WV. Repair of peripheral nerve defects in the upper extremity. *Hand Clin.* 2000 Feb;16(1):37-52.
2. Battiston B, Tos P, Conforti LG, Geuna S. Alternative techniques for peripheral nerve repair: conduits and end-to-side neuroorrhaphy. *Acta Neurochir Suppl.* 2007;100:43-50.
3. Oberlin C, Béal D, Leechavengvongs S, Salon A, Dauge MC, Sarcy JJ. Nerve transfer to biceps muscle using a part of ulnar nerve for C5-C6 avulsion of the brachial plexus: anatomical study and report of four cases. *J Hand Surg [Am].* 1994 Mar;19(2):232-7.
4. Loy S, Bhatia A, Asfazadourian H, Oberlin C. Ulnar nerve fascicle transfer onto to the biceps muscle nerve in C5-C6 or C5-C6-C7 avulsions of the brachial plexus. Eighteen cases]. *Ann Chir Main Memb Super.* 1997;16(4):275-84.
5. Teboul F, Kakkar R, Ameer N, Beaulieu JY, Oberlin C. Transfer of fascicles from the ulnar nerve to the nerve to the biceps in the treatment of upper brachial plexus palsy. *J Bone Joint Surg Am.* 2004 Jul;86-A(7):1485-90.
6. Mackinnon SE. Preliminary results of double nerve transfer to restore elbow flexion in upper type brachial plexus palsies. *Plast Reconstr Surg.* 2006 Oct;118(5):1273; author reply 1274.
7. Colbert SH, Mackinnon S. Posterior approach for double nerve transfer for restoration of shoulder function in upper brachial plexus palsy. *Hand (N Y).* 2006 Dec;1(2):71-7.
8. Mackinnon SE, Novak CB, Myckatyn TM, Tung TH. Results of reinnervation of the biceps and brachialis muscles with a double fascicular transfer for elbow flexion. *J Hand Surg [Am].* 2005 Sep;30(5):978-85.
9. Carlsen BT, Bishop AT, Shin AY. "Late reconstruction for brachial

plexus injury." Neurosurg Clin N Am. 2009 Jan;20(1):51-64, vi.

10. Cardenas-Mejia A, O'Boyle CP, Chen K, Chuang DC. Evaluation of Single-, Double-, and Triple-Nerve Transfers for Shoulder Abduction in 90 Patients with Supraclavicular Brachial Plexus Injury. *Plastic and Reconstructive Surgery.* Vol 2008 Nov;122(5):1470-1478.
11. Ranalli NJ, Kline DG, McGarvey ML, Boulis NM, Zager EL. Clinical Problem-Solving: Brachial Plexus Closed Injury and Reconstruction. *Neurosurgery* Vol 2008 Jun;62(6):1330-1339.
12. Goubier J, Teboul F. Technique of the Double Nerve Transfer to Recover Elbow Flexion in C5, C6, or C5 to C7 Brachial Plexus Palsy. *Techniques in Hand & Upper Extremity Surgery.* 2007 March;11(1):15-17.
13. Bertelli JA, Ghizoni MF. Results of grafting the anterior and posterior divisions of the upper trunk in complete palsies of the brachial plexus. *J Hand Surg [Am].* 2008 Nov;33(9):1529-40.
14. Bertelli JA, Santos MA, Kechele PR, Ghizoni MF, Duarte H. Triceps motor nerve branches as a donor or receiver in nerve transfers. *Neurosurgery.* 2007 Nov;61(5 Suppl 2):333-8; discussion 338-9.
15. Bertelli JA, Ghizoni MF. Transfer of the accessory nerve to the suprascapular nerve in brachial plexus reconstruction. *J Hand Surg [Am].* 2007 Sep;32(7):989-98.
16. Battiston B, Lanzetta M. Reconstruction of high ulnar nerve lesions by distal double median to ulnar nerve transfer. *J Hand Surg [Am].* 1999 Nov;24(6):1185-91.

Adham Abdelfattah, B.S., Paul L. Foster School of Medicine, El Paso, Texas.

Miguel Pirela-Cruz, M.D., Professor, Regional Chairman, Paul L. Foster School of Medicine, El Paso, Texas.

If you have recently changed your email address please notify the El Paso County Medical Society or you can log into the TMA's web page: www.texmed.org and make any corrections to your profile.